

# SHOPPERS DRUG MART. ...helping you save money

**PAMPERS DISPOSABLE DIAPERS**  
Toddler 48's  
Extra Absorbent 60's

**829**  
each



**Pampers**  
48's

600g  
**COMET**  
CLEANSER

**69¢**  
each



1 litre  
**MR. CLEAN**  
CLEANER

**179**  
each



750 ml  
**SCOPE**  
MOUTHWASH

**249**  
each



pkg. 4's  
**IVORY PERSONAL**  
BARS

**97¢**  
pkg.




150 ml  
**CREST**  
TOOTHPASTE

**169**  
each




**TOP NATIONAL BRANDS AT SHOPPERS DRUG MART MONEY SAVING PRICES**



## POISON remedy chart

Pick up your free poison remedy chart at the check out counter. This handy reference chart has self adhesive strips for fastening in a convenient location i.e. inside medicine cabinet, over sink in kitchen, by the telephone.



SHOPPERS DRUG MART

## POISON remedy chart

TYPE OF POISONING	WHAT TO DO
<b>SWALLOWED POISONS</b>	If the person is awake and able to swallow, give milk or water only. Then call the poison center or doctor. Caution: Antidote labels on products and antidote charts may be out of date and incorrect. <b>DO NOT</b> give salt, vinegar, or citrus fruit juices.
<b>POISONS ON THE SKIN</b>	Remove any affected clothing. Flood involved parts with water, wash with soap and water, and rinse. Then call the poison center or doctor.
<b>POISONS IN THE EYE</b>	Flood the eye with lukewarm (never hot) water, poured from a pitcher held 3-4 inches from the eye for 15 minutes. Then call the poison center or MD.
<b>INHALED POISONS</b>	Immediately, carry or drag the person to fresh air and give mouth-to-mouth resuscitation if necessary. Ventilate the area. Then call poison center or MD.

Always keep on hand copies of books, which include vomiting, and poison pills, which act as a laxative. Do not use either one unless instructed to do so by the poison center or your doctor, and follow their directions for use.

SOME COMMON SUBSTANCES & POSSIBLE SYMPTOMS OF OVERDOSE		
<b>ACETAMINOPHEN</b> Symptoms: Nausea, vomiting, loss of appetite, abdominal pain, jaundice (yellowing of the skin and eyes), dark urine, clay-colored stools, confusion, coma. Treatment: Supportive care, may require hospitalization.	<b>ASPIRIN</b> Symptoms: Nausea, vomiting, ringing in the ears, dizziness, confusion, tinnitus, bleeding gums, easy bruising, dark stools, bleeding from the stomach or intestines. Treatment: Supportive care, may require hospitalization.	<b>IBUPROFEN</b> Symptoms: Nausea, vomiting, dizziness, headache, blurred vision, ringing in the ears, confusion, tinnitus, bleeding gums, easy bruising, dark stools, bleeding from the stomach or intestines. Treatment: Supportive care, may require hospitalization.
<b>VALIUM</b> Symptoms: Drowsiness, slurred speech, impaired coordination, loss of consciousness, respiratory depression, hypotension, hypoxia, coma. Treatment: Supportive care, may require hospitalization.	<b>PHENYLEPHRINE</b> Symptoms: Rapid heart rate, high blood pressure, headache, dizziness, blurred vision, difficulty swallowing, difficulty breathing. Treatment: Supportive care, may require hospitalization.	<b>AMPHETAMINE</b> Symptoms: Rapid heart rate, high blood pressure, dilated pupils, dry mouth, loss of appetite, insomnia, tremor, sweating, hyperreflexia, muscle twitching, hallucinations, delirium, coma. Treatment: Supportive care, may require hospitalization.

EMERGENCY PHONE NUMBERS	
INDIAN CAPITAL CENTER	POLICE
FIRE DEPT.	HOSPITAL
EMERGENCY	AMBULANCE
	FIRE

Prices in effect until May 9th, 1981. We reserve the right to limit quantities while quantities last.