

# Soils and Goods

Address communications to Agronomist, 73 Adelaide St. West, Toronto.

## HANDLING THE YOUNG LAMBS.

Our experience has shown that the ewes that are in the best flesh and condition in the fall are the ones that breed first and, what is more essential, breed a larger per cent. of twins. We have heard flock masters say they "would rather have one good single lamb than twins." With any well-cared-for flock this is a great mistake. Our experience has shown for many years that we get far greater returns from the ewes having twins.

Now, for the care of the ewe and young lamb. First, provide a clean, dry place for the ewe to lamb. While a warm place is certainly desirable, yet we feel a clean, dry place means more than the warmth.

The wise shepherd will pass through his flock very often and note each individual carefully as lambing time comes on. A wrong presentation is not unusual. A normal presentation is the front feet with the nose and head immediately following. Sometimes the hind feet appear first in this case. Be sure the bottoms of the feet are turned up. Should they be otherwise, the shepherd, whose hands have first been carefully washed and disinfected, and with finger nails closely cut, and having thoroughly greased the hand with some soft, clean grease, should carefully insert the hand and using plenty of time get the fetus turned over and probably all will be well.

We have had the front feet come first O.K., but the head turned back. This simply means untold suffering and final death to both ewe and offspring unless assistance is given. In this case, after proper preparation as above described, the shepherd should take plenty of time and with great care, force back the fetus and following up until it is well back and while the head is yet there carefully bring the head forward and usually the trouble is over. Sometimes, but very rarely, the rump may try to come first and the ewe labor for hours and even days with no visible presentation and finally die. The remedy is obvious. Insert the hand, force back the fetus and secure a normal presentation either way.

Nearly all farmers, we feel sure, are in far too great haste to help, as they think, in the delivery. First, be sure the presentation is normal and then go away and leave nature alone for at least three or four hours. The parts need plenty of time to expand and yield sufficiently for the delivery, and great injury and even death may result from too great haste.

If present when the lambs are dropped, avoid handling as much as possible, but be sure they are properly cleaned up and dried. Next examine the ewe's udder and see the milk starts readily from both teats. A very little stoppage in the end of the teat will discourage the weak lamb and result in much trouble.

It may, and with many farmers does often occur in cold weather that the new born lamb becomes chilled and dies. In the case of chilled lambs, we use the warm water treatment. We found this treatment almost by accident many years ago and have had some most remarkable experiences with it, completely restoring lambs we had thought dead. We will not stop to relate them here, but for the benefit of some poor beginner will describe the process.

## Distention of Rumen.

Every winter many good animals are lost and many more nearly lost through carelessness in the tie-up, or the leaving of feed bins open. If a cow gets loose in the stable, she generally manages to find the bin where the feed grains are stored and just naturally gorges herself on the concentrated food. Can't blame the cow for eating. She knows not what trouble she is wading into by overloading her paunch with mill chaff or grain meal.

If it is known that an animal has gorged itself, and it can generally be determined by pressing the flank with the closed fist and noting if the indent of the hand remains for a short time, a drench should be given at once. One and a half pounds of Epsom salts dissolved in two gallons of water should be given at a single dose and followed by kneading the left side below the midian line vigorously to separate the compacted mass of fermenting grain and allow the drench to penetrate it. Stimulants, such as aromatic spirits of ammonia, should be given in two-ounce doses every four hours. Should the drench fail and the impacted condition continues, it can be relieved by a trained veterinarian performing the operation known as rumenotomy. This operation is performed by making an incision through the left flank and into the rumen, large enough to permit the entrance of the hand. The compacted grain or meal is removed in part (about two-thirds) and the wound in the rumen sutured up with catgut, after proper antiseptic precautions have been taken. The wound through the muscle and skin can be brought together with silk stitches, placed one

inch apart and through the entire thickness of the muscle. The wound should be dressed daily and kept covered to protect it from insects and dust. A useful dressing lotion can be made by using zinc sulphate, 1 dram; camphor acid, 2 drams; glycerine, 2 ounces; water, 14 ounces; mixed together. Should this operation be attempted by anyone other than a trained veterinary surgeon, warning is given that only an antiseptic, absolutely clean operation will permit the animal to live.

## In-Breeding and Line Breeding.

Having located a successful mating, one can perpetuate easily so long as the originals are breeding, but the problem begins after the first mating ceases. Shall one breed mother and son, or brother and sister, or cousins, or what is the best procedure?

Our experience, so says Professor W. R. Graham of the Ontario Agricultural College, in breeding birds of very close relationship, such as mother-son or brother-sister matings, would not warrant us recommending such a procedure. It is true that we have had some successful results, but it is likewise true that most of such matings have been very bad. If one is anxious to breed closely, our experience would suggest a trial of any relationship, so long as the birds are vigorous, but it is not wise to depend entirely upon such a mating. A good in-bred bird will likely be a valuable breeder, but such a bird is difficult to produce.

It is very doubtful if one can say what relationship will give the best results, but to the average person it would appear to be good advice not to breed too closely, and to look for new blood from some source where the breeding has been similar.

## The Construction of the Hotbed.

Many of the most popular vegetables, such as tomatoes, eggplant, peppers, celery, and of our best flowers, are of southern origin and require a longer growing season than this climate permits. By starting such plants in the hotbed from six to eight weeks earlier than it is safe to plant out doors, and later transplanting to the open, the required growing season may be obtained. Furthermore, by the use of the hotbed, such vegetables as radish, lettuce, cabbage and cauliflower may be started early and thus reach development much sooner than if the seed were sown in the open.

The only really satisfactory location for a hotbed is one with a southern exposure, protected from north and west winds and unshaded to admit a maximum amount of sunlight. It may be made either above or below ground as desired. The former requires more manure, the latter more labor to construct. If above ground, the bed should be situated where there is no danger of water standing at the time; if below, the situation should be naturally well drained, or drainage would have to be provided. With this latter type, the soil should be removed to a depth of about eighteen inches and the cavity should be at least two feet wider and longer than the frame to be used. This operation is best done during the previous autumn, as it is often quite practical to start a hotbed before the frost is out of the ground to a depth of 18 inches. With the above-ground type the site merely requires levelling, which may be done in the spring as well as any time.

The frame may be made of inch and a half or two inch lumber. The most convenient size is six feet wide and of the required length to use one or more sash three feet wide. It should be from ten to twelve inches high at the front and, if six feet wide, sixteen to eighteen at the back. This slope allows rain water to run off easily and as the hotbed should face south, a greater degree of sunlight is obtained than if the top were level. The sash must fit snugly and the frames be so constructed that they can be raised or lowered as ventilation is needed.

Fresh horse manure, as free from straw as possible, is the only satisfactory kind, for a hotbed, and that from grain fed animals is preferable. For a bed six feet square, one and a half to two tons is required. It should be secured and piled near the site a week or more before it is to be used, and turned as soon as it begins to heat, thus ensuring even heating and uniform texture. Four or five days after turning, the manure should be evenly heated and ready for the hotbed.

The bed of manure should be at least one foot wider every way than the frame to be placed upon it. It is laid down evenly in layers about six inches deep and each layer thoroughly tramped. The depth of manure required depends upon when the hotbed is made. If made early, two to two and a half feet are necessary, but if made during the latter part of March, one foot will be found sufficient. After manure has been placed to the proper depth the frame should be put on and more manure banked around the outside of it up to the top and at least one foot wide. The lights should

be applied to the animal.

Keep house plants clean and free from dust. Water thoroughly when the plants need water rather than a little every day.

When fed to horses in large quantities, wheat is apt to develop digestive disturbances and also cause skin eruptions. The best plan to follow in giving the grain to horses is to feed in moderate quantities only.

Grind it coarsely and mix with some other bulk feed like bran or chopped hay or straw.

## The New Wheat Champion

Canada has secured the world wheat championship for the twelfth time in her thirteenth year of endeavor. This year a new name has been added to the notable list of internationally famous grain growers who have gone one better than any one else, that of Major H. G. L. Strange, of Fenn, Alberta. The new name must also be added to that scarcely smaller list of inexperienced city men who in the Canadian West have outstripped farmers of lifelong labors and arrived at the pinnacle of agricultural fame, proving yet again that inexperience and lack of training constitute no handicap to successful farming in the Canadian West. Provided a man brings the right human qualities to bear upon his task and enthusiastically cooperates with Nature, who is so kindly disposed to that area. A civil engineer becomes added to the list already containing a tailor, a market gardener, and a book-stall clerk.

Major Strange, from the outset of his endeavors, has had only one object in view—the achievement of the best possible in all the lines of agricultural production he followed. He has his own implements for the production of special first generation seeds. With a 22-inch separator he threshes out small lots at the rate of about a sackful at a time. He is very much interested in hogs and has imported from England a number of the "large black" variety, which he claims was the original bacon type. At present he is believed to be the only breeder of this type of hog in Canada. He is an equal enthusiast in poultry, and trap-nests every one of his birds. All started home, happy to be together again.

A satisfactory winter ration for dairy cows kept at the Ste. Anne de la Pocatiere, Quebec, Experimental Station, consisted of corn and sunflower silage, roots, and mixed hay.

To this was added a meal mixture of oats, corn, bran, distillers' grains, and oil cake. This ration was fed at the rate of one pound of the mixture for every four pounds of milk produced by the individual cow.

DEVOTES TIME TO PUBLIC AFFAIRS.

The pursuit of agricultural perfection has not engrossed Major Strange to the exclusion of devoting his ability to public service, and, on the contrary, his visions in such work better and more profitable farming. He is the president of the Alberta Seed Growers' Association and the representative for Western Canada on the Dominion Advisory Seed Board. He is at the same time the secretary for the Alberta Poultry Breeders' Association, of which body he is a very active member.

It is no mere coincidence that so many city-trained men rise to the greatest heights of agricultural honor in Western Canada. The land of the Prairie Provinces is virgin, fertile and full of promise to be made productive to the extent of the human endeavor and intelligence applied. The entire farming history of Western Canada goes to prove that no inexperienced man need fear taking on the responsibilities of a farm in that territory, granted he brings intelligence, labor, and the right disposition to bear upon his task.

DECIDED TO SETTLE IN CANADA.

On leaving the army and facing the common problem of demobilized men, of where to go and the best course to pursue in the necessity of taking up the threads of life in a sadly disarranged world, Major Strange decided for the Canadian West and for mixed farming. He secured a farm in the famed Central Alberta region and concentrated on the production of elite seed and pure-bred livestock and pou-

now be placed on the frame and carefully closed so that the heat will be retained and the bed thus brought to a high temperature as soon as possible. After about two days the sash should be removed and the manure tramped thoroughly and watered if necessary. The hotbed is now ready for the soil. This should be of good fertility and of such a texture that it will not bake. If seed is to be sown directly in the hotbed, five or six inches of soil will be required, but if seed flats are to be used, two inches will be sufficient. No seed should be sown until the temperature in the hotbed has fallen to between eighty and ninety degrees Fahrenheit. After sowing, the temperature must be carefully observed, and if it gets too high the lights should be raised to provide ventilation. When the plants appear the frame should be kept aired sufficiently to prevent weak, spindly growth, although the plants must not be chilled or frozen. Water should be applied carefully, as too much will make for damping-off conditions. The soil, however, should be kept damp and this, together with sufficient air, light, and proper heat, will promote satisfactory growth.

## Chronic Tympanitis.

This form of bloat is present in many cattle stables during the long winter period when cattle are standing in. Some animals may be affected by this form of digestive trouble,

while others living under identical conditions are not.

The affected animal bloats up after feeding, but generally not to an extent to require puncture and relief by trocar and canula. A purgative will generally give relief, provided the chronic bloating has been caused by indigestion, and not by tuberculosis of the pulmonary lymphatic glands.

The following dosage has been found very effective in relieving this form of bloat: one pound of Epsom salts, one-half ounce of powdered Barbados aloes, one ounce of powdered ginger, and one pint of molasses. Stir all for five minutes and give as a drench. After the operation of the purgative see that the feed is sufficiently succulent to favor easy digestion.

Roots, silage and clover are most useful in keeping the alimentary tract in good condition. The use of linseed meal and gauze salts mixed with the feed morning and night—a handful of each—is good practice.

Should there be any indication that the general feeding is not laxative enough, potassium bi-carbonate, powdered ginger and powdered gentian are very useful as a tonic to tone up the digestive system and thereby aid in the prevention of chronic bloat or tympanitis.

Four equal parts of each, mix well and give one ounce at the beginning of the feed three times a day.

Should chronic bloating continue after treatment the tuberculin test should

be applied to the animal.

Keep house plants clean and free

from dust. Water thoroughly when the plants need water rather than a little every day.

When fed to horses in large quantities, wheat is apt to develop digestive disturbances and also cause skin eruptions. The best plan to follow in giving the grain to horses is to feed in moderate quantities only.

Grind it coarsely and mix with some other bulk feed like bran or chopped hay or straw.

## THE CHILDREN'S HOUR

### A HAPPY REUNION.

"Right, ho! We must not stand here idly. Let's get busy and become real detectives," said Rolly Rabbit, as soon as they found Bruin, gone from the trap. "Who will help me?"

"I—I will—I," chorused Frankie Fox, Willie Woodchuck and the little squirrel.

"We will let wise Frankie Fox be our chief," said Rolly. "He is always so clever at finding out secrets."

After Frankie had looked everything over carefully and had made several trips out among the trees, he said: "It is just as Rolly Rabbit thought first. Bruin went in that direction," pointing off toward the east. "His tracks seem to follow a man's and he must have been led away with a rope. We will follow them until we find him, and then plan some way of getting him away from the man. I am sure Bruin did not go away without being forced."

"So am I," chirped the little squirrel. "He was dreadfully afraid the man would come before we got back, and he did. But I ran just as fast as I could."

"You did your best, and if we are careful we may find Bruin before it gets dark," answered Rolly.

So off they started. Frankie Fox first, then Rolly Rabbit, then Willie Woodchuck, and trotting along behind was the little squirrel. Every few minutes Frankie would sniff the earth and nod his head, but they never stopped.

After a while it became dark and they could go no further. So they all curled up in a little nest of leaves by a big tree and went to sleep. Early in the morning they started on their way again. But Frankie Fox had slept so hard he just couldn't remember how Bruin's tracks smelled, and soon they were off the trail and lost.

All day long they wandered among the trees, and it was a weary and discouraged group that curled up on a bed of leaves that night.

Bright and early the next morning just as the sun was getting up, the little squirrel got up too.

"I'm so hungry, I must find some berries," he said. "I'll slip out quietly and not get far away."

Nearby he found a fine patch of blackberries and when busily munching them right in the middle of the patch, he stepped on something that caught his foot and made him fall.

"Ouchie, owl!" said the little squirrel, rubbing his bruised knee. "What was that? Why? It is Bruin's handkerchief, for it has his name on it."

Scampering back to his friends, he told them of his find and soon they were all on the right trail again. It was just an hour before they came upon Bruin sleeping by a log.

"Buzz-zz-z! Buzz-zz!" He was snoring so he never heard them.

Frankie Fox got a long blade of grass and tickled his ear, and Willie Woodchuck tickled his nose with a little twig. But when the squirrel tickled his toe, he jumped up with a start.

"Well, where did you all come from?" blinked Bruin, rubbing his eyes.

Then Rolly told him all about the dreadful time they had finding him, and Bruin told them about the man and how the little boy had unfastened his chain. With his sharp little knife, Rolly cut off Bruin's muzzle and they all started home, happy to be together again.

Back to Pre-War Prices

This splendid halter, in 1-1/2 inch clear red cedar, double door, double bars, 1-1/2 inch, double and stitched.

If your dealer cannot supply you, we will send it anywhere post paid, on receipt of price.


Price includes Shank.

100 Chick Brooder with Incubator \$19.75

100 Hen Capacity Grain Spreuer \$16.50

My twenty-four years experience has proved that the above prices are the lowest in the country.

L. R. Guild, Incubator Dept., Rockwood, Ont.



## Live Stock in Canada.

Dominion-wide statistics indicate that there were fewer horses, mules, cattle, and sheep in Canada at the end of 1923 than in 1922 and more swine and poultry. The figures supplied are: Horses, in 1923, 8,530,641 compared with 8,648,871 in 1922, a decrease of 118,230; mules, 8,722 compared with 9,202, a decrease of 480; cattle, 9,246,281 compared with 9,719,869, a decrease of 473,638; sheep, 2,753,860 compared with 3,263,525, a decrease of 509,665; swine, 4,405,316 compared with 3,915,634, an increase of 489,632; poultry, 45,469,292 compared with 42,930,562, an increase of 2,538,750. Horses and sheep have decreased in every province excepting British Columbia; cattle have increased in Ontario and British Columbia but have decreased in every other province; swine have decreased in New Brunswick and Nova Scotia; but have increased in all the other provinces; poultry have increased in Ontario, Saskatchewan, Alberta, and British Columbia but have decreased in the Maritime Provinces, Quebec and Manitoba. These figures do not include the live stock on Indian reserves.

A recent survey of the hog population of Prussia shows a substantial increase, the number now on hand totaling 9,460,000, as compared with 8,683,936 a year ago.

To be a good farmer you must feed your mind, your body, your soil, and your soul. They are merely engines that will run if you give them enough of the right kind of fuel, and will stop and rust if you don't.

## BABY CHICKS